

UTHOU-16UTL 79-88.ST25 SUPPLEMENTAL SEQUENCE LISTING

```
. <110> The Board of Regents of the University of Texas System
 <120>
        MUTATIONS IN A NOVEL PHOTORECEPTOR-PINEAL GENE ON 17P CAUSE LEBER
         CONGENITAL AMAUROSIS (LCA4)
 <130>
        96606/16UTL
 <140>
                                                                       RECEIVED
        09/765,061
 <141>
        2001-01-17
 <150>
        60/331362
                                                                          JUN 1 1 2003
 <151>
        2001-01-04
 <160>
                                                                     TECH CENTER 1600/2900
        10 additional sequences, Seq. Nos. 79-88
 <170> PatentIn version 3.2
 <210>
        79
 <211>
        34
 <212>
        DNA
 <213>
        Homo sapiens
 <220>
 <221>
        misc_feature
 <222>
        (1)..(34)
 <223>
        Donor Splice Site: Residue 1-10 are the exonic sequence and
        Residues 11-34 are the intronic sequence
 <400>
 cggatcccga gtgagtgggg ccctccggag caga
                                                                            34
 <210>
        80
 <211>
        35
 <212>
        DNA
 <213>
        Homo sapiens
                                                                                1.
 <220>
 <221>
<222>
        misc_feature
        (1)..(35)
 <223>
        Acceptor Splice Site: Residues 1-25 are the intronic sequence and Residues 26-35 are the exonic sequence.
 <400>
 cagagtgcac cgtctcggtg actaggtgat ctttc
                                                                            35
 <210>
        81
 <211>
        35
 <212>
        DNA
 <213>
        Homo sapiens
 <220>
 <221>
        misc_feature
 <222>
        (1)..(35)
 <223>
        Donor Splice Site: Residue 1-10 are the exonic sequence and
        Residues 11-35 are the intronic sequence
```

UTHou-16UTL 79-88.ST25 csacaccatc gtaagtaggc cctgcgcgcc tgtct 35 82 35 <210> <211> <212> DNA <213> Homo sapiens <220> <221> misc_feature <222> (1)..(35)<223> Acceptor Splice Site: Residues 1-25 are the intronic sequence and Residues 26-35 are the exonic sequence. <400> gccatccatc cgtttatccc cacagcacac ggggg 35 <210> 83 <211> 35 <212> DNA <213> Homo sapiens <220> <221> misc_feature <222> (1)..(35)<223> Donor Splice Site: Residue 1-10 are the exonic sequence and Residues 11-35 are the intronic sequence <400> 83 gctgctgcag gtggggctgg ggttggcagg gctgg 35 <210> 84 <211> 35 <212> DNA <213> Homo sapiens <220> <221> <222> misc_feature (1)..(35)<223> Acceptor Splice Site: Residues 1-25 are the intronic sequence and Residues 26-35 are the exonic sequence. cactgacctg cagctctggg gccaggttga tgccc 35 <210> 85 <211> 35 <212> DNA <213> Homo sapiens <220> misc_feature <221> <222> (1)..(35)Donor Splice Site: Residue 1-10 are the exonic sequence and <223> Residues 11-35 are the intronic sequence

Page 2

UTHou-16UTL 79-88.ST25

٠	<400> gcagaco	85 caag gtcagaggcc gctggccacg gggtg	35
r	<210> <211> <212> <213>	86 35 DNA Homo sapiens	
	<220> <221> <222> <223>	misc_feature (1)(35) Acceptor Splice Site: Residues 1-25 are the intronic sequence and Residues 26-35 are the exonic sequence.	
	<400> catggc	86 tgac cttctccctg ggcaggagaa gccrt	35
	<210> <211> <212> <213>	87 35 DNA Homo sapiens	
2	<220> <221> <222> <223>	misc_feature (1)(35) Donor Splice Site: Residue 1-10 are the exonic sequence and Residues 11-35 are the intronic sequence	
	<400> caccaco	87 ccag gtgcgcgggg ctgcaggggc ggaca	35
	<210> <211> <212> <213>	88 35 DNA Homo sapiens	
	<220> <221> <222> <223>	misc_feature (1)(35) Acceptor Splice Site: Residues 1-25 are the intronic sequence and Residues 26-35 are the exonic sequence.	
	<400>	88' Tact contactor canadicate atasa	35